

p-ニトロアニソールのラットを用いた経口投与による
2 週 間 毒 性 試 験（混 餌 試 験）報 告 書

試験番号：0360

TABLES

TABLES

TABLE 1	EXPERIMENTAL DESIGN AND MATERIALS AND METHODS IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE
TABLE 2	SURVIVAL ANIMAL NUMBERS AND BODY WEIGHT CHANGES OF MALE RATS IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE
TABLE 3	SURVIVAL ANIMAL NUMBERS AND BODY WEIGHT CHANGES OF FEMALE RATS IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE
TABLE 4	FOOD CONSUMPTION CHANGES OF MALE RATS IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE
TABLE 5	FOOD CONSUMPTION CHANGES OF FEMALE RATS IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE

TABLE 1 EXPERIMENTAL DESIGN AND MATERIALS AND METHODS
IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE

2-week study	
<Method of Administration>	Mixed feed
<Number of Groups>	Male 6, Female 6
<Size of Groups>	5 males and 5 females of each group
<Animals>	
Strain and Species	F344/DuCrj (Fischer) rat
Animal Source	Charles River Japan, Inc.
Duration Held Before Study	2 wk
Age When Placed on Study	6 wk
Age When Killed	8 wk
<Doses>	
<Male>	0, 1250, 2500, 5000, 10000 or 20000 ppm
<Female>	0, 1250, 2500, 5000, 10000 or 20000 ppm
<Duration of Dosing>	7d/wk for 2wk
<Animal Maintenance>	
Feed	CRF-1 (Oriental Yeast Co., Ltd.) Sterilized by γ -ray Available <i>ad libitum</i>
Water	Filtrated and sterilized by ultraviolet ray Automatic watering system Available <i>ad libitum</i>
Animal per Cage	Single (stainless steel wire)
Animal Room Environment	Barrier system Temperature : $23\pm 2^{\circ}\text{C}$ Humidity : $55\pm 15\%$ Fluorescent light 12h/d 15~17 room air changes /h
<Type and Frequency of Observation>	
Clinical Sign	Observed 0-0, 1-1, 1-3, 1-7, 2-3, and 2-7 (wk-d)
Body Weight	Weighed 0-0, 1-1, 1-3, 1-7, 2-3, and 2-7 (wk-d)
Food Consumption	Weighed 1-3, 1-7, 2-3, and 2-7 (wk-d)

TABLE 1 EXPERIMENTAL DESIGN AND MATERIALS AND METHODS
(Continued) IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE

2-week study	
<Hematology>	Red blood cell (RBC), Hemoglobin, Methemoglobin, Hematocrit, Mean corpuscular volume (MCV), Mean corpuscular hemoglobin (MCH), Mean corpuscular hemoglobin concentrate (MCHC), Platelet, Reticulocyte, White blood cell (WBC), Differential WBC, Prothrombin time (PT), Activated partial thromboplastin time (APTT).
<Biochemistry>	Total protein, Albumin, A/G ratio, Total bilirubin, Glucose, Total cholesterol Phospholipid, Glutamic oxaloacetic transaminase (GOT), Glutamic pyruvic transaminase (GPT), Lactate dehydrogenase (LDH), γ -Glutamyl transpeptidase (G-GTP), Creatine phosphokinase (CPK), Urea nitrogen, Creatinine, Sodium, Potassium, Chloride, Calcium, Inorganic phosphorus.
<Urinalysis>	None
<Necropsy>	Necropsy performed on all animals.
<Organ Weight>	Organ weight measurement performed on at least five animals per sex per group at sacrifice The following organs were weighed; thymus, adrenal, testis, ovary, heart, lung, kidney, spleen, liver, and brain.
<Histopathologic Examination>	Histopathologic examination performed on at least two animals per sex per group. The following organs were examined; skin, nasal cavity, nasopharynx, larynx, trachea, lung, bone marrow, lymph node, thymus, spleen, heart, tongue, salivary gland, esophagus, stomach, small intestine, large intestine, liver, pancreas, kidney, urinary bladder, pituitary, thyroid, parathyroid, adrenal, testis, epididymis, seminal vesicle, prostate, ovary, uterus, vagina, mammary gland, brain, spinal cord, peripheral nerve, eye, Harderian gland, muscle, bone, other organs/tissues with gross lesions.

TABLE 2 SURVIVAL ANIMAL NUMBERS AND BODY WEIGHT CHANGES OF MALE RATS
IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE

Week-day on Study	Control			1250ppm			2500ppm			5000ppm			10000ppm			20000ppm		
	Av.Wt.	No. of Surviv. <5>		Av.Wt.	% of cont. <5>	No. of Surviv.	Av.Wt.	% of cont. <5>	No. of Surviv.	Av.Wt.	% of cont. <5>	No. of Surviv.	Av.Wt.	% of cont. <5>	No. of Surviv.	Av.Wt.	% of cont. <5>	No. of Surviv.
0-0	130	(5)	5/5	129	(5)	99	130	(5)	100	129	(5)	99	129	(5)	99	129	(5)	99
1-1	136	(5)	5/5	133	(5)	98	131	(5)	96	125	(5)	92	120	(5)	88	116	(5)	85
1-3	144	(5)	5/5	142	(5)	99	141	(5)	98	134	(5)	93	120	(5)	83	108	(5)	75
1-7	161	(5)	5/5	160	(5)	99	159	(5)	99	153	(5)	95	135	(5)	84	102	(5)	63
2-3	178	(5)	5/5	176	(5)	99	174	(5)	98	168	(5)	94	145	(5)	81	99	(5)	56
2-7	192	(5)	5/5	190	(5)	99	190	(5)	99	184	(5)	96	157	(5)	82	87	(5)	45
< > : No.of effective animals, () : No.of measured animals Av.Wt.:g																		

TABLE 3 SURVIVAL ANIMAL NUMBERS AND BODY WEIGHT CHANGES OF FEMALE RATS
IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE

Week-day on Study	Control			1250ppm			2500ppm			5000ppm			10000ppm			20000ppm		
	Av.Wt.	No. of Surviv. <5>		Av.Wt.	% of cont. <5>	No. of Surviv.	Av.Wt.	% of cont. <5>	No. of Surviv.	Av.Wt.	% of cont. <5>	No. of Surviv.	Av.Wt.	% of cont. <5>	No. of Surviv.	Av.Wt.	% of cont. <5>	No. of Surviv.
0-0	98	(5)	5/5	98	(5)	100	98	(5)	100	98	(5)	100	98	(5)	100	98	(5)	100
1-1	99	(5)	5/5	100	(5)	101	97	(5)	98	93	(5)	94	90	(5)	91	88	(5)	89
1-3	103	(5)	5/5	103	(5)	100	102	(5)	99	97	(5)	94	88	(5)	85	79	(5)	77
1-7	111	(5)	5/5	111	(5)	100	109	(5)	98	105	(5)	95	95	(5)	86	69	(5)	62
2-3	119	(5)	5/5	118	(5)	99	112	(5)	94	112	(5)	94	98	(5)	82	61	(4)	51
2-7	126	(5)	5/5	123	(5)	98	120	(5)	95	118	(5)	94	107	(5)	85	-	(0)	-
< > : No.of effective animals, () : No.of measured animals Av.Wt.:g																		

TABLE 4 FOOD CONSUMPTION CHANGES OF MALE RATS IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE

Week-day on Study	Control			1250ppm			2500ppm			5000ppm			10000ppm			20000ppm							
	Av.FC.	No. of Surviv.		Av.FC.	% of cont.	No. of Surviv.	Av.FC.	% of cont.	No. of Surviv.	Av.FC.	% of cont.	No. of Surviv.	Av.FC.	% of cont.	No. of Surviv.	Av.FC.	% of cont.	No. of Surviv.					
		<5>			<5>			<5>			<5>			<5>			<5>						
1-3	13.8	(5)	5/5	12.9	(5)	93	5/5	11.8	(5)	86	5/5	8.8	(5)	64	5/5	6.3	(5)	46	5/5	5.5	(5)	40	5/5
1-7	14.0	(5)	5/5	13.4	(5)	96	5/5	13.0	(5)	93	5/5	12.0	(5)	86	5/5	9.7	(5)	69	5/5	6.3	(5)	45	5/5
2-3	14.7	(5)	5/5	14.2	(5)	97	5/5	14.1	(5)	96	5/5	13.6	(5)	93	5/5	10.5	(5)	71	5/5	6.9	(5)	47	5/5
2-7	13.9	(5)	5/5	14.1	(5)	101	5/5	14.2	(5)	102	5/5	14.1	(5)	101	5/5	11.5	(5)	83	5/5	7.8	(5)	56	5/5
< > : No.of effective animals, () : No.of measured animals Av.FC.:g																							

TABLE 5 FOOD CONSUMPTION CHANGES OF FEMALE RATS IN THE 2-WEEK FEED STUDY OF p-NITROANISOLE

Week-day on Study	Control			1250ppm				2500ppm				5000ppm				10000ppm				20000ppm			
	Av.FC.		No. of Surviv.	Av.FC.		% of cont.	No. of Surviv.	Av.FC.		% of cont.	No. of Surviv.	Av.FC.		% of cont.	No. of Surviv.	Av.FC.		% of cont.	No. of Surviv.	Av.FC.		% of cont.	No. of Surviv.
			<5>			<5>			<5>			<5>			<5>			<5>			<5>		
1-3	9.7	(5)	5/5	9.2	(5)	95	5/5	8.5	(5)	88	5/5	7.2	(5)	74	5/5	8.5	(5)	88	5/5	8.5	(5)	88	5/5
1-7	10.2	(5)	5/5	9.6	(5)	94	5/5	9.1	(5)	89	5/5	8.7	(5)	85	5/5	6.7	(5)	66	5/5	6.2	(5)	61	5/5
2-3	10.7	(5)	5/5	9.7	(5)	91	5/5	8.9	(5)	83	5/5	9.2	(5)	86	5/5	6.8	(5)	64	5/5	9.6	(4)	90	4/5
2-7	10.1	(5)	5/5	9.7	(5)	96	5/5	9.4	(5)	93	5/5	9.2	(5)	91	5/5	7.8	(5)	77	5/5	-	(0)	-	0/5
< > : No.of effective animals, () : No.of measured animals Av.FC.:g																							