

4-*tert*-ブチルカテコールのマウスを用いた
経口投与による 13 週間毒性試験（混餌試験）報告書

試験番号：0720

APPENDICES

APPENDICES

- APPENDIX 1-1 IDENTITY OF 4-*tert*-BUTYLCATECHOL IN THE 13-WEEK FEED STUDY
- APPENDIX 1-2 STABILITY OF 4-*tert*-BUTYLCATECHOL IN THE 13-WEEK FEED STUDY
- APPENDIX 2-1 CONCENTRATION OF 4-*tert*-BUTYLCATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY
- APPENDIX 2-2 HOMOGENEITY OF 4-*tert*-BUTYLCATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY
- APPENDIX 2-3 STABILITY OF 4-*tert*-BUTYLCATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY
- APPENDIX 3 METHODS, UNITS AND DECIMAL PLACE FOR HEMATOLOGY AND BIOCHEMISTRY IN THE 13-WEEK FEED STUDY OF 4-*tert*-BUTYLCATECHOL

APPENDIX 1-1


IDENTITY OF 4-*tert*-BUTYLCATECHOL IN THE 13-WEEK FEED STUDY

Infrared Spectrometry


Instrument : Shimadzu FTIR-8200PC Infrared Spectrometer

Cell : KBr

Resolution : 4 cm^{-1}



Infrared Spectrum of Test Substance



Infrared Spectrum of Literature Data*

Result: The infrared spectrum was consistent with literature spectrum.
(*Performed by Wako Pure Chemical Industries, Ltd.)

2. Conclusion: The test substance was identified as 4-*tert*-butylcatechol by mass spectrum and infrared spectrum.

IDENTITY OF 4-*tert*-BUTYLcatechol IN THE 13-WEEK FEED STUDYTest Substance : 4-*tert*-Butylcatechol (Wako Pure Chemical Industries, Ltd.)

Lot No. : PEK3567



1. Spectral Data

Mass Spectrometry

Instrument : Hitachi M-80B Mass Spectrometer

Ionization : EI (Electron Ionization)

Ionization Voltage : 70eV



Result: The mass spectrum was consistent with literature spectrum.

(*McLafferty FW, ed. 1994. Wiley Registry of Mass Spectral Data. 6th ed. New York, NY:John Wiley and Sons.)

APPENDIX 1-2

STABILITY OF 4-*tert*-BUTYLCATECHOL IN THE 13-WEEK FEED STUDY

STABILITY OF 4-*tert*-BUTYLCATECHOL IN THE 13-WEEK FEED STUDYTest Substance : 4-*tert*-Butylcatechol (Wako Pure Chemical Industries, Ltd.)

Lot No. : PEK3567

1. High Performance Liquid Chromatography

Instrument : Shimadzu LC-10 High Performance Liquid Chromatograph

Column : TSK-GEL ODS-80TM (4.6 mm ϕ $\times$ 15 cm)

Column Temperature: 40 °C

Flow Rate : 1 mL/min

Mobile Phase : Acetonitrile : 5mmol SDS solution (Phosphoric acid pH2.2) = 50 : 50

Detector : UV (285 nm)

Injection Volume : 10 μ L

Date analyzed	Peak No.	Retention Time (min)	Area (%)
2008.10.31	1	4.299	100
2009.02.18	1	4.301	100

Result: High performance liquid chromatography indicated one major peak (peak No.1) analyzed on 2008.10.31 and one major peak (peak No.1) analyzed on 2009.2.18. No new trace impurity peak in the test substance analyzed on 2009.2.18 was detected.

2. Conclusion: The test substance was stable for the period that the test substance had been used for the study.

APPENDIX 2-1

CONCENTRATION OF 4-*tert*-BUTYLCATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

CONCENTRATION OF 4-*tert*-BUTYL-CATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

Analytical Method : The samples were analyzed by high performance liquid chromatography.

Instrument : Shimadzu LC-10 High Performance Liquid Chromatograph

Column : TSK-GEL ODS-80TM (4.6 mm ϕ $\times$ 15 cm)

Column Temperature: 40 °C

Flow Rate : 1 mL/min

Mobile Phase : Acetonitrile : 5mmol SDS solution (Phosphoric acid pH2.2) = 50 : 50

Detector : UV (285 nm)

Injection Volume : 10 μ L

Date Analyzed	Target Concentration				
	625 ^a	1250	2500	5000	10000
2008.11.12	657 ^b (105) ^c	1340 (107)	2680 (107)	5100 (102)	9580 (95.8)

^a ppm

^b ppm (Mean measured concentration.)

^c % (Mean measured concentration/target concentration $\times$ 100.)

APPENDIX 2-2

HOMOGENEITY OF 4-*tert*-BUTYLCATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

HOMOGENEITY OF 4-*tert*-BUTYLCATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

Analytical Method : The samples were analyzed by high performance liquid chromatography.
 Instrument : Shimadzu LC-10 High Performance Liquid Chromatograph
 Column : TSK-GEL ODS-80TM (4.6 mm ϕ $\times$ 15 cm)
 Column Temperature: 40 °C
 Flow Rate : 1 mL/min
 Mobile Phase : Acetonitrile : 5mmol SDS solution (Phosphoric acid pH2.2) = 50 : 50
 Detector : UV (285 nm)
 Injection Volume : 10 μ L

	Target Concentration				
	625 ^a	1250	2500	5000	10000
Coefficient Variation	2.12 ^b	2.01	1.89	2.37	3.17

^a ppm^b % (n=7)

APPENDIX 2-3

STABILITY OF 4-*tert*-BUTYLCATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

STABILITY OF 4-*tert*-BUTYLCATECHOL IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

Analytical Method : The samples were analyzed by high performance liquid chromatography.

Instrument : Shimadzu LC-10 High Performance Liquid Chromatograph

Column : TSK-GEL ODS-80TM (4.6 mm ϕ $\times$ 15 cm)

Column Temperature: 40 °C

Flow Rate : 1 mL/min

Mobile Phase : Acetonitrile : 5mmol SDS solution (Phosphoric acid pH2.2) = 50 : 50

Detector : UV (285 nm)

Injection Volume : 10 μ L

Date Analyzed	Target Concentration	
	150 ^a	15000
2008.10.28	144 (100) ^b	14200 (100)
2008.11.05 ^c	134 (93.1)	13700 (96.5)
2008.11.05 ^d	139 (96.5)	13800 (97.2)

^a ppm

^b % (Percentage was based on the concentration at the date of preparation.)

^c Animal room samples

^d Cold storage samples

APPENDIX 3

METHODS, UNITS AND DECIMAL PLACE FOR HEMATOLOGY AND BIOCHEMISTRY IN THE 13-WEEK FEED STUDY OF 4-*tert*-BUTYLCATECHOL

**METHODS, UNITS AND DECIMAL PLACE FOR HEMATOLOGY AND BIOCHEMISTRY
IN THE 13- WEEK FEED STUDY STUDY OF 4-*tert*-BUTYLCA TECHOL**

Item	Method	Unit	Decimal place
Hematology			
Red blood cell (RBC)	Light scattering method ¹⁾	$\times 10^6/\mu\text{L}$	2
Hemoglobin(Hgb)	Cyanmethemoglobin method ¹⁾	g/dL	1
Hematocrit(Hct)	Calculated as $\text{RBC} \times \text{MCV}/10$ ¹⁾	%	1
Mean corpuscular volume(MCV)	Light scattering method ¹⁾	fL	1
Mean corpuscular hemoglobin(MCH)	Calculated as $\text{Hgb}/\text{RBC} \times 10$ ¹⁾	pg	1
Mean corpuscular hemoglobin concentration (MCHC)	Calculated as $\text{Hgb}/\text{Hct} \times 100$ ¹⁾	g/dL	1
Platelet	Light scattering method ¹⁾	$\times 10^3/\mu\text{L}$	0
Reticulocyte	Light scattering method ¹⁾	%	1
Methemoglobin	Van Assendelft method ²⁾	%	1
White blood cell(WBC)	Light scattering method ¹⁾	$\times 10^3/\mu\text{L}$	2
Differential WBC	Light scattering method ¹⁾	%	0
Biochemistry			
Total protein(TP)	Biuret method ³⁾	g/dL	1
Albumin (Alb)	BCG method ³⁾	g/dL	1
A/G ratio	Calculated as $\text{Alb}/(\text{TP} - \text{Alb})$ ³⁾	—	1
T-bilirubin	Azobilirubin method ³⁾	mg/dL	2
Glucose	GlcK·G-6-PDH method ³⁾	mg/dL	0
T-cholesterol	CE·COD·POD method ³⁾	mg/dL	0
Triglyceride	MGLP·GK·GPO·POD method ³⁾	mg/dL	0
Phospholipid	PLD·ChOD·POD method ³⁾	mg/dL	0
Aspartate aminotransferase (AST)	JSCC method ³⁾	IU/L	0
Alanine aminotransferase (ALT)	JSCC method ³⁾	IU/L	0
Lactate dehydrogenase (LDH)	JSCC method ³⁾	IU/L	0
Alkaline phosphatase (ALP)	JSCC method ³⁾	IU/L	0
γ -Glutamyl transpeptidase (γ -GTP)	JSCC method ³⁾	IU/L	0
Creatine kinase (CK)	JSCC method ³⁾	IU/L	0
Urea nitrogen	Urease·GLDH method ³⁾	mg/dL	1
Sodium	Ion selective electrode method ³⁾	mEq/L	0
Potassium	Ion selective electrode method ³⁾	mEq/L	1
Chloride	Ion selective electrode method ³⁾	mEq/L	0
Calcium	OCPC method ³⁾	mg/dL	1
Inorganic phosphorus	PNP·XOD·POD method ³⁾	mg/dL	1

1) Automatic blood cell analyzer (ADVIA120 : Siemens Healthcare Diagnostics Inc.)

2) Spectrophotometer (DU-530 : Beckman Coulter, Inc.)

3) Automatic analyzer (Hitachi 7080 : Hitachi, Ltd.)