4-クロロ-2-ニトロアニリンのラットを用いた 経口投与による13週間毒性試験(混餌試験)報告書

試験番号:0745

APPENDICES

APPENDICES

- APPENDIX 1-1 IDENTITY OF 4-CHLORO-2-NITROANILINE IN THE 13-WEEK FEED STUDY
- APPENDIX 1-2 STABILITY OF 4-CHLORO-2-NITROANILINE IN THE 13-WEEK FEED STUDY
- APPENDIX 2-1 CONCENTRATION OF 4-CHLORO-2-NITROANILINE IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY
- APPENDIX 2-2 HOMOGENEITY OF 4-CHLORO-2-NITROANILINE IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY
- APPENDIX 2-3 STABILITY OF 4-CHLORO-2-NITROANILINE IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY
- APPENDIX 3 METHODS, UNITS AND DECIMAL PLACE FOR HEMATOLOGY AND BIOCHEMISTRY IN THE 13-WEEK FEED STUDY OF 4-CHLORO-2-NITROANILINE

APPENDIX 1-1

IDENTITY OF 4-CHLORO-2-NITROANILINE

IN THE 13-WEEK FEED STUDY

IDENTITY OF 4-CHLORO-2-NITROANILINE IN THE 13-WEEK FEED STUDY

Test Substance : 4-Chloro-2-nitroaniline (Tokyo Chemical Industry Co., Ltd.)


Lot No. : GJ01


1. Spectral Data

Mass Spectrometry


- Instrument : Hitachi M-80B Mass Spectrometer
- Ionization : EI (Electron Ionization)

Ionization Voltage : 70eV






Infrared Spectrometry




Resolution : 4 cm^{-1}

)



Infrared Spectrum of Literature Data*



2. Conclusion: The test substance was identified as 4-chloro-2-nitroaniline by mass spectrum and infrared spectrum.

APPENDIX 1-2

STABILITY OF 4-CHLORO-2-NITROANILINE

IN THE 13-WEEK FEED STUDY

STABILITY OF 4-CHLORO-2-NITROANILINE IN THE 13-WEEK FEED STUDY

Test Substance : 4-Chloro-2-nitroaniline (Tokyo Chemical Industry Co., Ltd.)

Lot No. : GJ01

1. High Performance Liquid Chromatography

Instrument : Shimadzu LC-10 High Performance Liquid Chromatograph

Column : TSK-GEL ODS-80TM (4.6 mm $\phi \times 15$ cm)

Column Temperature: 40 °C

Flow Rate : 1 mL/min

Mobile Phase : Acetonitrile : 5mmol SDS solution (Phospholic acid pH2.2) = 70 : 30

Detector : UV (405 nm)

Injection Volume : 10 µL

ļ

Date analyzed	Peak No.	Retention Time (min)	Area (%)
2009.10.09	1	3.047	100
2010.02.01	1	3.046	100

- Result: High performance liquid chromatography indicated one major peak (peak No.1) analyzed on 2009.10.9 and one major peak (peak No.1) analyzed on 2010.2.1. No new trace impurity peak in the test substance analyzed on 2010.2.1 was detected.
- 2. Conclusion: The test substance was stable for the period that the test substance had been used for the study.

APPENDIX 2-1

CONCENTRATION OF 4-CHLORO-2-NITROANILINE IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

CONCENTRATION OF 4-CHLORO-2-NITROANILINE IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

Analytical Method	: The samples were analyzed by high performance liquid chromatography.
Instrument	: Shimadzu LC-10 High Performance Liquid Chromatograph
Column	: TSK-GEL ODS-80TM (4.6 mm ϕ $ imes$ 15 cm)
Column Temperatu	are: 40 °C
Flow Rate	: 1 mL/min
Mobile Phase	: Acetonitrile : 5mmol SDS solution (Phospholic acid pH2.2) = 70 : 30
Detector	: UV (405 nm)
Injection Volume	: 10 μL

	Target Concentration				
Date Analyzed	640ª	1600	4000	7000	10000
2009.10.12	697 ^b (109) ^c	1730 (108)	4340 (109)	7510 (107)	10700 (107)

^a ppm

^b ppm (Mean measured concentration.)
^c % (Mean measured concentration/target concentration × 100.)

APPENDIX 2-2

HOMOGENEITY OF 4-CHLORO-2-NITROANILINE IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

HOMOGENEITY OF 4-CHLORO-2-NITROANILINE IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

Analytical Method	: The samples were analyzed by high performance liquid chromatography.
Instrument	: Shimadzu LC-10 High Performance Liquid Chromatograph
Column	: TSK-GEL ODS-80TM (4.6 mm ϕ $ imes$ 15 cm)
Column Temperatu	re: 40 °C
Flow Rate	: 1 mL/min
Mobile Phase	: Acetonitrile : 5mmol SDS solution (Phospholic acid pH2.2) = 70 : 30
Detector	: UV (405 nm)
Injection Volume	: 10 µL

	Target Concentration				
	640ª	1600	4000	7000	10000
Coefficient Variation	6.91 ^b	6.52	2.09	2.70	4.37

^a ppm

^b % (n=7)

APPENDIX 2-3

STABILITY OF 4-CHLORO-2-NITROANILINE

IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

STABILITY OF 4-CHLORO-2-NITROANILINE IN FORMULATED DIETS IN THE 13-WEEK FEED STUDY

Analytical Method	: The samples were analyzed by high performance liquid chromatography.
Instrument	: Shimadzu LC-10 High Performance Liquid Chromatograph
Column	: TSK-GEL ODS-80TM (4.6 mm ϕ $ imes$ 15 cm)
Column Temperatu	re: 40 °C
Flow Rate	: 1 mL/min
Mobile Phase	: Acetonitrile : 5mmol SDS solution (Phospholic acid pH2.2) = 70 : 30
Detector	: UV (405 nm)
Injection Volume	: 10 µL

	Target Concentration			
Date Analyzed	640ª	10000	100	
2009.09.09	612 (100) ^b	9970 (100)	101 (100)	
2009.09.17°	579 (94.6)	9670 (97.0)	94.4 (93.5)	
2009.09.17 ^d	645 (105)	10500 (105)	95.3 (94.4)	

^a ppm

)

^b % (Percentage was based on the concentration at the date of preparation.)
 ^c Animal room samples
 ^d Cold storage samples

APPENDIX 3

METHODS, UNITS AND DECIMAL PLACE FOR HEMATOLOGY AND BIOCHEMISTRY IN THE 13-WEEK FEED STUDY OF 4-CHLORO-2-NITROANILINE

METHODS, UNITS AND DECIMAL PLACE FOR HEMATOLOGY AND BIOCHEMISTRY IN THE 13- WEEK FEED STUDY STUDY OF 4-CHLORO-2-NITROANILINE

Item	Method	Unit	Decimal	
			place	
Hematology	× ·			
Red blood cell (RBC)	Light scattering method ¹⁾	$ imes$ 106/ μ L	2	
Hemoglobin(Hgb)	Cyanmethemoglobin method ¹⁾	g/dL	1	
Hematocrit(Hct)	Calculated as RBC \times MCV/10 ¹⁾	%	1	
Mean corpuscular volume(MCV)	Light scattering method ¹⁾	fL	1	
Mean corpuscular hemoglobin(MCH)	Calculated as Hgb/RBC \times 10 ¹⁾	pg	1	
Mean corpuscular hemoglobin concentration	Calculated as Hgb/Hct $ imes$ 100 $^{1)}$	g/dL	1	
(MCHC)				
Platelet	Light scattering method ¹⁾	$ imes 10^{3}/\mu\mathrm{L}$	0	
Reticulocyte	Light scattering method ¹⁾	%	1	
Methemoglobin	Van Assendelft method ²⁾	%	1	
White blood cell(WBC)	Light scattering method ¹⁾	$ imes 10^{3/} \mu { m L}$	2	
Differential WBC	Light scattering method ¹⁾	%	0	
Biochemistry				
Total protein(TP)	Biuret method ³⁾	g/dL	1	
Albumin (Alb)	BCG method ³⁾	g/dL	1	
A/G ratio	Calculated as Alb/(TP-Alb) ³⁾	_	1	
T-bilirubin	Azobilirubin method ³⁾	mg/dL	2	
Glucose	GlcK·G-6-PDH method ³⁾	mg/dL	0	
T -cholesterol	CE·COD·POD method ³⁾	mg/dL	0	
Triglyceride	MGLP·GK·GPO·POD method ³⁾	mg/dL	0	
Phospholipid	PLD·ChOD·POD method ³⁾	mg/dL	0	
Aspartate aminotransferase (AST)	JSCC method ³⁾	IU/L	0	
Alanine aminotransferase (ALT)	JSCC method ³⁾	IU/L	0	
Lactate dehydrogenase (LDH)	JSCC method ³⁾	IU/L	0	
Alkaline phosphatase (ALP)	JSCC method ³⁾	IU/L	0	
γ -Glutamyl transpeptidase (γ -GTP)	JSCC method ³⁾	IU/L	0	
Creatine kinase (CK)	JSCC method ³⁾	IU/L	0	
Urea nitrogen	Urease · GLDH method ³⁾	mg/dL	1	
Creatinine	Jaffé method ³⁾	mg/dL	1	
Sodium	Ion selective electrode method ³⁾	mÉq/L	0	
Potassium	Ion selective electrode method ³⁾	mEq/L	1	
Chloride	Ion selective electrode method ³⁾	mEq/L	0	
Calcium	OCPC method ³⁾	mg/dL	1	
Inorganic phosphorus	PNP·XOD·POD method ³⁾	mg/dL	1	

1) Automatic blood cell analyzer (ADVIA120 : Siemens Healthcare Diagnostics Inc.)

2) Spectrophotometer (DU-530 : Beckman Coulter, Inc.)

3) Automatic analyzer (Hitachi 7080 : Hitachi, Ltd.)

)

)